



## Microplastic exposure studies should be environmentally realistic

Lenz, Robin; Enders, Kristina; Nielsen, Torkel Gissel

*Published in:*  
Proceedings of the National Academy of Sciences

*Link to article, DOI:*  
[10.1073/pnas.1606615113](https://doi.org/10.1073/pnas.1606615113)

*Publication date:*  
2016

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Lenz, R., Enders, K., & Nielsen, T. G. (2016). Microplastic exposure studies should be environmentally realistic. *Proceedings of the National Academy of Sciences*, 113(29), E4121 - E4122 . [201606615].  
<https://doi.org/10.1073/pnas.1606615113>

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Microplastic exposure studies should be environmentally realistic

Robin Lenz<sup>a,1</sup>, Kristina Enders<sup>a</sup>, and Torkel Gissel Nielsen<sup>a</sup>

To understand the impact of microplastic (MP) pollution to aquatic ecosystems, it is important to identify the mechanisms of interaction with organisms. Exposure experiments, like the study of Sussarellu et al. (1) recently published in PNAS, may provide such insights. However, the results of dose–response experiments must always be interpreted in light of environmental concentrations, and the experimental concentrations examined by Sussarellu et al. (1) and several others (2–8) are orders-of-magnitude higher than those reported from field studies (Fig. 1).

Experimental studies on effects of MP on mussels (2, 3), lugworms (4), copepods (5–7), and oysters (1) have documented reduced feeding, survival, and fecundity, as well as promoted polychlorinated biphenyl bioaccumulation linked to MP uptake. In addition, nano-sized plastic particles may reduce body size and lead to reproductive malfunctioning in *Daphnia* (8), and to reduced CO<sub>2</sub> uptake and enhanced production of reactive oxygen species in algae (9).

All of these experimental studies use MP concentrations far above the levels documented in the marine environment (Fig. 1). Many of those studies used MP beads smaller than those reported from the field, but even if one extrapolates field-measured concentrations of MP particles to the size range used in experiments, experimental and expected field concentrations are very far apart. Studies that determined concentrations for micro-bead exposures from field measurements of larger-sized MP assumed a mass-conserving fragmentation

tation process, where numbers of particles will scale inversely with the particle radius to the power of 3. In contrast, the environmental concentrations documented in studies appear to follow a slightly lower exponent (2.67) (Fig. 1), possibly caused by size-dependent removal processes, lower dimensional breakdown (i.e., of flakes, sheets, and fibers), or considerable influence of new MP input of larger sizes. Additionally, decreased detection accuracy of applied sampling methods in their respective lower size ranges might play a role.

Experimental exposure concentrations tend to be between two to seven orders-of-magnitude higher than environmental levels. The most recent study (1) used concentrations closest to those found in nature. However, Sussarellu et al. calculate their MP concentrations based on sediment data from a pollution hot-spot area close to a ship-breaking yard in India (8, 10); thus, these data are unlikely to be representative of general concentrations beyond the local area.

Microplastic research is an emerging field, and there is a lot of misunderstanding and in some cases overreaction or misinterpretation of results from MP science in the public. We therefore strongly suggest that future studies of MP impact on marine ecosystems should also include concentrations that have been documented in the environment to yield more realistic estimates of sublethal effects.

## Acknowledgments

The authors thank Thomas Kiørboe and Mary Wisz for helpful comments on this letter.

<sup>a</sup>Section for Marine Ecology and Oceanography, National Institute of Aquatic Resources, Technical University of Denmark, 2920 Charlottenlund, Denmark  
 Author contributions: K.E. and T.G.N. designed research; R.L. performed research; T.G.N. provided scientific leadership; and R.L. and K.E. wrote the paper.

The authors declare no conflict of interest.

Data deposition: A fully referenced version of the data review has been deposited at [figshare.com](https://figshare.com), accessible under: [https://figshare.com/articles/Size\\_and\\_concentration\\_comparison\\_for\\_microplastics\\_between\\_reported\\_environmental\\_levels\\_and\\_laboratory\\_exposure\\_studies/3206449](https://figshare.com/articles/Size_and_concentration_comparison_for_microplastics_between_reported_environmental_levels_and_laboratory_exposure_studies/3206449).

<sup>1</sup>To whom correspondence should be addressed. Email: [roble@aqua.dtu.dk](mailto:roble@aqua.dtu.dk).

